



Towards a Dynamical Systems Approach to Mental Disorders

Adela Isvoranu

Department of Psychological Research Methods
University of Amsterdam

Part 1. Story of Origins

Psychology as a Complex System




Cramer, A. O. J., Waldorp, L., van der Maas, H., & Borsboom, D. (2010). Comorbidity: A Network Perspective. *Behavioral and Brain Sciences*, 33(2-3), 137– 150. doi: 10.1017/S0140525X09991567

Borsboom, D. (2017). A network theory of mental disorders. *World Psychiatry*, 16(1):5–13.

Psychology as a Complex System

- Psychological processes are increasingly understood as **complex dynamical systems** of interacting components
 - ✓ Psychological
 - ✓ Biological
 - ✓ Sociological
 - ✓ ...
- Recent literature (~10 years) in the field of mental disorders started embracing this complexity

Psychology as a Complex System




Cramer, A. O. J., Waldorp, L., van der Maas, H., & Borsboom, D. (2010). Comorbidity: A Network Perspective. *Behavioral and Brain Sciences*, 33(2-3), 137–150. doi: 10.1017/S0140525X09991567


Borsboom, D. (2017). A network theory of mental disorders. *World Psychiatry*, 16(1):5–13.

Analogy from Ecology

General factor



Mutualism




van der Maas, H. L., Dolan, C. V., Grasman, R. P., Wicherts, J. M., Huizenga, H. M., & Raijmakers, M. E. (2006). A dynamical model of general intelligence: The positive manifold of intelligence by mutualism. *Psychological Review*, 113(4), 842–861.
doi: 10.1037/0033-295X.113.4.842

Network Psychometrics

Psychological networks differ from networks of many other disciplines in one very fundamental aspect

- Edges are *not observed* and need to be *estimated*




Epskamp, S., Borsboom, D., & Fried, E. I. (2018). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1), 195-212.

Network Psychometrics

Psychological networks differ from networks of many other disciplines in one very fundamental aspect

- Edges are *not observed* and need to be *estimated*




Typically two steps:

- (1) Estimate a statistical model on data, from which some parameters can be represented as a weighted network between observed variables
- (2) Analyze the weighted network structure using measures taken from graph theory to infer, for instance, the most central nodes.

Epskamp, S., Borsboom, D., & Fried, E. I. (2018). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1), 195-212.

Network Psychometrics


- A node represents an entity
 - People
 - Cities
 - **Symptoms**
 - **Psychological constructs**












Network Psychometrics

An edge represents some connection between two nodes

- ❑ Friendship / contact
- ❑ Distance
- ❑ **Comorbidity**
- ❑ **Causality**
- ❑ **Interaction**




Network Psychometrics: Psycho(patho)logy



Papers using Network Psychometrics (up to June 2018):


- Anker, J. J., Forbes, M. K., Almquist, Z. W., Menk, J. S., Thuras, P., Unruh, A. S., & Kushner, M. G. (2017). A network approach to modeling comorbid internalizing and alcohol use disorders. *Journal of Abnormal Psychology*, 126(3), 325–339. <https://doi.org/10.1037/abn0000257>
- Armour, C., Fried, E. I., Deserno, M. K., Tsai, J., Pietrzak, R. H., Southwick, S. M., & Pietrzak, R. H. (2017). A network analysis of DSM-5 posttraumatic stress disorder symptoms and correlates in U.S. military veterans. *Journal of Anxiety Disorders*, 45, 49–59. Retrieved from <http://eiko-fried.com/wp-content/uploads/1-s2.0-S0887618516302419-main-1.pdf>
- Baggio, S., Starcevic, V., Studer, J., Simon, O., Gainsbury, S. M., Gmel, G., & Billieux, J. (n.d.). Technology-mediated addictive behaviors constitute a spectrum of related yet distinct conditions: A network perspective. Short title: Network of technology-mediated addictions. Retrieved from https://www.researchgate.net/profile/joel_Billieux/publication/324836030_Technology-mediated_addictive_behaviors_constitute_a_spectrum_of_related_yet_distinct_conditions_A_network_perspective/links/5ae6ffddaca272ba50821ec1/Technology-mediated-addictive-behaviors-constitute-a-spectrum-of-related-yet-distinct-conditions-A-network-perspective.pdf
- Bak, M., Drukker, M., Hasmi, L., & van Os, J. (2016). An n=1 Clinical Network Analysis of Symptoms and Treatment in Psychosis. *PLOS ONE*, 11(9), e0162811. <https://doi.org/10.1371/journal.pone.0162811>
- Beard, C., Millner, A. J., Forgeard, M. J. C., Fried, E. I., Hsu, K. J., Treadway, M. T., Fritz, J., Fried, E. I., Goodyer, I. M., Wilkinson, P. O., & Harmelen, A.-L. Van. (2016). Network analysis of depression and anxiety symptom relationships in a psychiatric sample. *Psychological Medicine*, 46(16), 3359–3369. <https://doi.org/10.1017/S0033291716002300>
- Bell, V., & O'Driscoll, C. (2018). The network structure of paranoia in the general population. *Social Psychiatry and Psychiatric Epidemiology*, 1–8. <https://doi.org/10.1007/s00127-018-1487-0>
- Bernstein, E. E., Heeren, A., & McNally, R. J. (2017). Unpacking Rumination and Executive Control: A Network Perspective. *Clinical Psychological Science*, 5(5), 816–826. <https://doi.org/10.1177/2167702617702717>
- Birkeland, M. S., & Heir, T. (2017). Making connections: exploring the centrality of posttraumatic stress symptoms and covariates after a terrorist attack. *European Journal of Psychotraumatology*, 8(sup3), 1333387. <https://doi.org/10.1080/20008198.2017.1333387>
- Blanken, T. F., Deserno, M. K., Dalege, J., Borsboom, D., Blanken, P., Kerkhof, G. A., & Cramer, A. O. J. (2018). The role of stabilizing and communicating symptoms given overlapping communities in psychopathology networks. *Scientific Reports*, 8(1), 5854. <https://doi.org/10.1038/s41598-018-24224-2>
- Bodner, N., Kuppens, P., Allen, N. B., Sheeber, L. B., & Ceulemans, E. (2017). Affective family interactions and their associations with adolescent depression: A dynamic network approach. *Development and Psychopathology*, 1–15. <https://doi.org/10.1017/S0954579417001699>
- Bos, F. M., Fried, E. I., Hollon, S. D., Bringmann, L. F., Dimidjian, S., DeRubeis, R. J., & Bockting, C. L. H. (2018). Cross-sectional networks of depressive symptoms before and after antidepressant medication treatment. *Social Psychiatry and Psychiatric Epidemiology*, 53(6), 617–627. <https://doi.org/10.1007/s00127-018-1506-1>
- Boschloo, L., Van Borkulo, C. D., Rhemtulla, M., Keyes, K. M., Borsboom, D., & Schoevers, R. A. (2015). The network structure of symptoms of the diagnostic and statistical manual of mental disorders. *PLOS ONE*, 10(9), e0137621. <https://doi.org/10.1371/journal.pone.0137621>
- Braginti, G., Kempenaers, C., Braun, S., Fried, E. I., & Linkowski, P. (2018). Network analysis of empathy items from the interpersonal reactivity index in 1973 young adults. *Psychiatry Research*, 265, 87–92. <https://doi.org/10.1016/J.PSYCHRES.2018.03.082>
- Bringmann, L. F., Lemmens, J. H. J. M., Huijbers, M. J. H., Borsboom, D., &
- Fisher, A. J., Reeves, J. W., Lawyer, G., Medaglia, J. D., & Rubel, J. A. (2017). Exploring the Idiographic Dynamics of Mood and Anxiety via Network Analysis. *Journal of Abnormal Psychology*, 126(8), 1044–1056. <https://doi.org/10.1037/abn0000311>
- Forrest, L. N., Jones, P. J., Ortiz, S. N., & Smith, A. R. (2018). Core psychopathology in anorexia nervosa and bulimia nervosa: A network analysis. *International Journal of Eating Disorders*. <https://doi.org/10.1002/eat.22871>
- Fried, E., Eidhof, M. B., Palic, S., Costantini, G., Huisman-van Dijk, H. M., Boekting, C. L. H., ... Karstoft, K.-I. (n.d.). Replicability and generalizability of PTSD networks: A cross-cultural multisite study of PTSD symptoms in four trauma patient samples. *Clinical Psychological Science*. <https://doi.org/10.17605/osf.io/2t7qp>
- Fried, E. I., Epskamp, S., Nesse, R. M., Tuerlinckx, F., & Borsboom, D. (2016). What are "good" depression symptoms? Comparing the centrality of DSM and non-DSM symptoms of depression in a network analysis. *Journal of Affective Disorders*, 189, 314–320. <https://doi.org/10.1016/j.jad.2016.01.030>
- Fried, E. I., van Borkulo, C. D., Cramer, A. O. J., Boschloo, L., Schoevers, R. A., & Borsboom, D. (2017). Mental disorders as networks of problems: a review of recent insights. *Social Psychiatry and Psychiatric Epidemiology*, 52(1), 1–10. <https://doi.org/10.1007/s00127-016-1319-z>
- Galderisi, S., Rucci, P., Kirkpatrick, B., Mucci, A., Gibertoni, D., Rocca, P., ... Maj, M. (2018). Interplay Among Psychopathologic Variables, Personal Resources, Context-Related Factors, and Real-life Functioning in Individuals With Schizophrenia. *JAMA Psychiatry*, 75(4), 396. <https://doi.org/10.1001/jamapsychiatry.2017.4607>
- Gittins Stone, D. I., DuBois, R., DeJesus, B., Rodgers, R. F., & Rizzo, C. (2017). Brief report: A network analysis of self-cutting risk among late adolescent girls exposed to dating violence. *Journal of Adolescence*, 61, 12–16. <https://doi.org/10.1016/j.adolescence.2017.09.002>
- Glück, T. M., Kneifel, M., & Lueger-Schuster, B. (2017). A network analysis of anger, shame, proposed ICD-11 post-traumatic stress disorder, and different types of childhood trauma in foster care settings in a sample of adult survivors. *European Journal of Psychotraumatology*, 8(sup3), 1372543. <https://doi.org/10.1080/20008198.2017.1372543>
- Goldschmidt, A. B., Crosby, R. D., Cao, L., Moessner, M., Forbush, K. T., Accurso, E. C., & Le Grange, D. (2018). Network analysis of pediatric eating disorder symptoms in a treatment-seeking, transdiagnostic sample. *Journal of Abnormal Psychology*, 127(2), 251–264. <https://doi.org/10.1037/abn0000327>
- Goold, C., Vas, J., Olsen, C., & Newberry, R. C. (2016). Using network analysis to study behavioural phenotypes: an example using domestic dogs. *Royal Society Open Science*, 3(10), 160268. <https://doi.org/10.1098/rsos.160268>
- Gouvernet, B., Combaluzier, S., Sebbe, F., & Rezrazi, A. (2017). Plurality and prevalence of sexual motivations in a sample of young francophone adults. *Revue Européenne de Psychologie Appliquée/European Review of Applied Psychology*, 67(5), 231–245. <https://doi.org/10.1016/j.erap.2017.08.002>
- Haslbeck, J. M. B., & Fried, E. I. (2017). How predictable are symptoms in psychopathological networks? A reanalysis of 18 published datasets. *Psychological Medicine*, 47(16), 2767–2776. <https://doi.org/10.1017/S0033291717001258>
- Hasson-Ohavon, I. (2018). Overlap and distinction between measures of insight obsessive-compulsive disorder and depression: a Bayesian network approach. *Psychological Medicine*, 47(7), 1204–1214. <https://doi.org/10.1017/S0033291716003287>
- Mcnally, R. J., Robinaugh, D. J., Wu, G. W. Y., Wang, L., Deserno, M. K., Borsboom, D., ... Borsboom, D. (2015). Mental disorders as causal systems: a network approach to posttraumatic stress disorder. *Clinical Psychological Science*, 3(6), 836–849. <https://doi.org/10.1037/cpsc0000025>
- Mitchell, K. S., Wolf, E. J., Bovin, M. J., Lee, L. O., Green, J. D., Rosen, R. C., ... Marx, B. P. (2017). Network models of DSM-5 posttraumatic stress disorder: Implications for ICD-11. *Journal of Abnormal Psychology*, 126(3), 355–366. <https://doi.org/10.1037/abn0000252>
- Mullarkey, M. C., Marchetti, I., & Beevers, C. G. (2018). Using Network Analysis to Identify Central Symptoms of Adolescent Depression. *Journal of Clinical Child & Adolescent Psychology*, 1–13. <https://doi.org/10.1080/15374416.2018.1437735>
- Murphy, J., McBride, O., Fried, E., & Shevlin, M. (2017). Distress, Impairment and the Extended Psychosis Phenotype: A Network Analysis of Psychotic Experiences in an US General Population Sample. *Schizophrenia Bulletin*. <https://doi.org/10.1093/schbul/sbx134>
- Núñez, D., Fresno, A., van Borkulo, C. D., Courtet, P., Arias, V., Garrido, V., & Wigman, J. T. W. (2018). Examining relationships between psychotic experiences and suicidal ideation in adolescents using a network approach. *Schizophrenia Research*. <https://doi.org/10.1016/j.schres.2018.05.020>
- Olatunji, B. O., Levinson, C., & Caleb, B. (2018). A network analysis of eating disorder symptoms and characteristics in an inpatient sample. *Psychiatry Research*, 262, 270–281. <https://doi.org/10.1016/J.PSYCHRES.2018.02.027>
- Pavani, J.-B., Le Vigouroux, S., Kop, J.-L., Congard, A., & Dauvier, B. (2017). A Network Approach to Affect Regulation Dynamics and Personality Trait-Induced Variations: Extraversion and Neuroticism Moderate Reciprocal Influences Between Affect and Affect Regulation Strategies. *European Journal of Personality*, 31(4), 329–346. <https://doi.org/10.1002/per.2109>
- Pereira-Morales, A. J., Adan, A., & Forero, D. A. (2017). Network analysis of multiple risk factors for mental health in young Colombian adults. *Journal of Mental Health*, 1–9. <https://doi.org/10.1080/09638237.2017.1417568>
- Preszler, J., Marcus, D. K., Edens, J. F., & McDermott, B. E. (2018). Network analysis of psychopathy in forensic patients. *Journal of Abnormal Psychology*, 127(2), 171–182. <https://doi.org/10.1037/abn0000325>
- Richetin, J., Preti, E., Costantini, G., & De Panfilis, C. (2017). The centrality of affective instability and identity in Borderline Personality Disorder: Evidence from network analysis. *PLOS ONE*, 12(10), e0186695. <https://doi.org/10.1371/journal.pone.0186695>
- Robinaugh, D. J., LeBlanc, N. J., Vuletic, H. A., & McNally, R. J. (2014). Network analysis of persistent complex bereavement disorder in conjugal bereaved adults. *Journal of Abnormal Psychology*, 123(3), 510–522. <https://doi.org/10.1037/abn0000002>
- Ross, J., Murphy, D., & Armour, C. (2018). A network analysis of DSM-5 posttraumatic stress disorder and functional impairment in UK treatment-seeking veterans. *Journal of Anxiety Disorders*. <https://doi.org/10.1016/j.janxdis.2018.05.007>
- Ruzzano, L., Borsboom, D., & Geurts, H. M. (2015). Repetitive Behaviors in Autism and Obsessive-Compulsive Disorder: New Perspectives from a Network Analysis. *Journal of Autism and Developmental Disorders*, 45(1), 192–202. <https://doi.org/10.1007/s10803-014-2204-9>
- Santos, H., Fried, E. I., Asufa-Adjei, J., & Ruiz, R. J. (2017). Network Structure of Perinatal Depressive Symptoms in Latinas: Relationship to Stress and

Network Psychometrics: Psycho(patho)logy




Depression

A




B




Fried, E. I., Epskamp, S., Nesse, R. M., Tuerlinckx, F., & Borsboom, D. (2016). What are 'good' depression symptoms? Comparing the centrality of DSM and non-DSM symptoms of depression in a network analysis. *Journal of affective disorders*, 189, 314-320.

Substance Use




Rhemtulla, M., Fried, E. I., Aggen, S. H., Tuerlinckx, F., Kendler, K. S., & Borsboom, D. (2016). Network analysis of substance abuse and dependence symptoms. *Drug and alcohol dependence*, 161, 230-237.

Psychosis




van Rooijen, G., **Ivoranu, A. M.**, Meijer, C. J., van Borkulo, C.D., Ruhé H. G., de Haan, L., GROUP (in press). A Symptom Network Structure of the Psychosis Spectrum. *Schizophrenia Research*.


Network Psychometrics: Psychopathology




External Influence on a Network Model




External Influence on a Network Model



External Influence on a Network Model




Environmental Impact in Psychotic Disorders



Ivoranu, A. M., Borsboom, D., van Os, J. & Guloksuz, S. (2016). A Network Approach to Environmental Impact in Psychotic Disorder: Brief Theoretical Framework. *Schizophrenia Bulletin*, 42(4), 870-873.


Environmental Impact in Psychotic Disorders




Ivoranu, A. M., Borsboom, D., van Os, J. & Guloksuz, S. (2016). A Network Approach to Environmental Impact in Psychotic Disorder: Brief Theoretical Framework. *Schizophrenia Bulletin*, 42(4), 870-873.

Environmental Impact in Psychotic Disorders

(b) No Environmental Exposure




(c) Environmental Exposure Cannabis




Ivoranu, A. M., Borsboom, D., van Os, J. & Guloksuz, S. (2016). A Network Approach to Environmental Impact in Psychotic Disorder: Brief Theoretical Framework. *Schizophrenia Bulletin*, 42(4), 870-873.

Pathways Between Childhood Trauma and Psychotic Symptoms

a. Shortest Paths: Sexual Abuse – Positive and Negative Symptoms





- Positive Symptoms
- Negative Symptoms
- GP Psychopathology
- Childhood Trauma



Ivoranu, A. M., van Borkulo, C. D., Boyette, L. L., Wigman, J. T. W., Vinkers, C. H., Borsboom, D. (2017). A Network Approach to Psychosis: Pathways Between Childhood Trauma and Psychotic Symptoms. Schizophrenia Bulletin, 43(1), 187-196.

Pathways Between Childhood Trauma and Psychotic Symptoms

a. Shortest Paths: Sexual Abuse – Positive and Negative Symptoms



Ivoranu, A. M., van Borkulo, C. D., Boyette, L. L., Wigman, J. T. W., Vinkers, C. H., Borsboom, D. (2017). A Network Approach to Psychosis: Pathways Between Childhood Trauma and Psychotic Symptoms. *Schizophrenia Bulletin*, 43(1), 187-196.

Part 3. Implications & Conclusions

Consequences & Conclusions

- ▣ Co-occurrence of symptoms, moods, personality aspects understood as emergent behavior
 - ▣ Implication: symptom-based interventions possible
- ▣ Every person is a different system
 - ▣ Implication: personalized modeling and treatment
- ▣ Possible to think in terms of attractor states with smooth or critical transitions
 - ▣ Implication: dynamical models may lead to insight in optimal intervention
- ▣ Possible to think about theory formation

Facebook Community



facebook.com/groups/PsychologicalDynamics

Thank you for your attention!



Adela Isvoranu

Twitter @Adelaisvoranu
www.adelaisvoranu.com

Facebook group: Psychological Dynamics

Slides at:
www.adelaisvoranu.com/presentations

